

Sets with two associative operations

By TEIMURAZ PIRASHVILI

*A.M. Razmadze Math. Inst. Aleksidze str. 1, Tbilisi,
380093. Republic of Georgia*

1 Introduction

Jean-Louis Loday introduced the notion of a *dimonoid* and a *dialgebra*. Let us recall, that a *dimonoid* is a set equipped with two associative operations satisfying 3 more axioms (see [2, 3] or Section 6.2), while a *dialgebra* is just a linear analog of a dimonoid.

In this paper we drop these additional axioms and we consider sets equipped with two associative binary operations. We call such an algebraic structure as a *duplex*. Thus dimonoids are examples of duplexes. The set of all permutations, gives an example of a duplex which is not a dimonoid (see Section 3). In Section 4 we construct a free duplex generated by a given set via planar trees and then we prove that the set of all permutations form a free duplex on an explicitly described set of generators. In the last section we consider duplexes coming from planar binary trees and vertices of the cubes as in [5]. We prove that these duplexes are free with one generator in appropriate variety of duplexes.

2 Graded sets

A *graded set* is a set X_* together with a decomposition $X_* = \coprod_{n \in \mathbb{N}} X_n$. Here and elsewhere $\coprod$ denotes the disjoint union of sets. A map of graded sets is a map $f : X_* \rightarrow Y_*$ such that $f(X_n) \subset Y_n$, $n \geq 0$. We let $\mathbf{Sets}_*$ be the category of graded sets. If $x \in X_n$ we write $\mathcal{O}(x) = n$. In this way we obtain a map $\mathcal{O} : X_* \rightarrow \mathbb{N}$ and conversely, if such a map is given then X_* is graded with $X_n = \mathcal{O}^{-1}(n)$. If X_* and Y_* are graded sets, then the Cartesian product $X_* \times Y_*$ is also a graded set with

$$\mathcal{O}(x, y) := \mathcal{O}(x) + \mathcal{O}(y), \quad x \in X_*, y \in Y_*.$$

The disjoint union of two graded sets X_* and Y_* is also graded by $(X_* \coprod Y_*)_n = X_n \coprod Y_n$. A graded set X_* is called *locally finite* if X_n is finite for any $n \in \mathbb{N}$. In this case we put

$$X_*(\mathbf{T}) := \sum_{n=0}^{\infty} \text{Card}(X_n) \mathbf{T}^n \in \mathbb{Z}[[\mathbf{T}]].$$

An n -ary operation $\flat$ on a graded set X_* is *homogeneous* if $\flat : X_*^n \rightarrow X_*$ is a map of graded sets, in other words

$$\flat(X_{k_1} \times \cdots \times X_{k_n}) \subset X_{k_1 + \cdots + k_n}.$$

Free semigroups are examples of graded sets. Let us recall that a *semigroup* is a set equipped with an associative binary operation. For any set S the free semigroup on S is the following graded set

$$\text{Ass}(S) = \coprod_{n \geq 1} \text{Ass}_n(S); \quad \text{Ass}_n(S) := S^n, n \geq 0$$

while the multiplication is given by

$$S^n \times S^m \rightarrow S^{n+m}; \quad ((x_1, \dots, x_n), (y_1, \dots, y_m)) \mapsto (x_1, \dots, x_n, y_1, \dots, y_m).$$

It is clear that

$$\text{Ass}(S)(\mathbf{T}) = \frac{\text{Card}(S)\mathbf{T}}{1 - \text{Card}(S)\mathbf{T}}.$$

3 Two operations on permutations

For any $n \geq 1$ we let $\underline{n}$ be the set $\{1, 2, \dots, n\}$. Furthermore, we let $\mathfrak{S}_n$ be the set of all bijections $\underline{n} \rightarrow \underline{n}$. An element $f \in \mathfrak{S}_n$ called a *permutation*, is specified by the sequence $(f(1), \dots, f(n))$. The composition law $\circ$ yields the group structure on $\mathfrak{S}_n$. We put

$$\mathfrak{S} := \coprod_{n=1}^{\infty} \mathfrak{S}_n.$$



We are going to consider $\mathfrak{S}$ as a graded set with respect of this grading. Thus $\mathcal{O}(f) = n$ means that $f : \underline{n} \rightarrow \underline{n}$ is a bijection. On $\mathfrak{S}$ we introduce two homogeneous associative operations. The first one is *the concatenation*, which we denote by $\sharp$. Thus

$$\sharp : \mathfrak{S}_n \times \mathfrak{S}_m \rightarrow \mathfrak{S}_{n+m},$$


is defined by

$$\begin{aligned} (f \sharp g)(i) &= f(i), & \text{if } 1 \leq i \leq n, \\ (f \sharp g)(i) &= n + g(i - n), & \text{if } n + 1 \leq i \leq n + m. \end{aligned}$$

Here $\mathcal{O}(f) = n$ and $\mathcal{O}(g) = m$. For example if $f \in \mathfrak{S}_3$ and $g \in \mathfrak{S}_3$ are the following permutations



then $f\sharp g \in \mathfrak{S}_6$ is given by




The second operation

$$\natural : \mathfrak{S}_n \times \mathfrak{S}_m \rightarrow \mathfrak{S}_{n+m},$$

is defined by

$$\begin{aligned} (f\natural g)(i) &= m + f(i), & \text{if } 1 \leq i \leq n, \\ (f\natural g)(i) &= g(i - n), & \text{if } n + 1 \leq i \leq n + m. \end{aligned}$$

For example if $f \in \mathfrak{S}_3$ and $g \in \mathfrak{S}_3$ are as above, then $f\natural g \in \mathfrak{S}_6$ is given by



These operations appears also in [5] under the name ‘over’ and ‘under’.

3.1 Duplexes

One easily checks that both operations $\sharp, \natural$ defined on $\mathfrak{S}$ are associative (see also Lemma 1.10 in [5]). This suggests to introduce the following definition

Definition 3.1 *A duplex is a set D equipped with two associative operations $\cdot : D \times D \rightarrow D$ and $*$: $D \times D \rightarrow D$. A map $f : D \rightarrow D'$ from a duplex D to another duplex D' is a homomorphism, provided $f(x \cdot y) = f(x) \cdot f(y)$ and $f(x * y) = f(x) * f(y)$.*

We let **Duplexes** be the category of duplexes. As usual one can introduce the notion of *free duplex* as follows.

Definition 3.2 *A duplex F is called free if there exists a subset $X \subset F$, such that for any duplex D and any map $f : X \rightarrow D$ there exists an unique homomorphism of duplexes $g : F \rightarrow D$ such that $g(x) = f(x)$ for all $x \in X$. If this holds, then we say that F is a free duplex on X .*

Thus, duplexes generalize the notion of dimonoids introduced by Jean-Louis Loday in [3]. Our main result is the following

Theorem 3.3 *The set $\mathfrak{S}$ equipped with $\sharp$ and $\natural$ is a free duplex on the set $\mathcal{U}^{\mathfrak{S}_2}$.*

The description of the set $\mathcal{U}^{\mathfrak{S}_2}$ and the proof of Theorem 3.3 is given in Section 5.

3.2 Semigroups $(\mathfrak{S}, \sharp)$ and $(\mathfrak{S}, \natural)$

In this section we prove the fact that the semi-group $(\mathfrak{S}, \sharp)$ is free. It is easy to see that $(\mathfrak{S}, \sharp)$ and $(\mathfrak{S}, \natural)$ are isomorphic semi-groups (see Lemma 3.4) and therefore $(\mathfrak{S}, \natural)$ is free as well.

Lemma 3.4 *Let $\omega_n \in \mathfrak{S}_n$ be the permutation defined by*

$$\omega_n(i) = n + 1 - i, \quad i \in \underline{n}$$

and let $\xi_n : \mathfrak{S}_n \rightarrow \mathfrak{S}_n$ be the map given by $\xi_n(f) = \omega_n \circ f$. Then

$$\xi_n \circ \xi_n = Id_{\mathfrak{S}_n}$$

and the diagram

$$\begin{array}{ccc} \mathfrak{S}_n \times \mathfrak{S}_m & \xrightarrow{\sharp} & \mathfrak{S}_{n+m} \\ \xi_n \times \xi_m \downarrow & & \downarrow \xi_{n+m} \\ \mathfrak{S}_n \times \mathfrak{S}_m & \xrightarrow{\natural} & \mathfrak{S}_{n+m} \end{array}$$

commutes. Therefore the map $\xi = \coprod_n \xi_n : \mathfrak{S} \rightarrow \mathfrak{S}$ is an isomorphism from the semi-group $(\mathfrak{S}, \sharp)$ to the semi-group $(\mathfrak{S}, \natural)$.

Proof. Since $\omega \circ \omega = Id_{\underline{n}}$ it follows that $\xi_n(\xi_n(f)) = \omega_n \circ (\omega_n \circ f) = f$. Furthermore, for any $f \in \mathfrak{S}_n$, $g \in \mathfrak{S}_m$ and $1 \leq i \leq n + m$, both $\xi_{n+m}(f \sharp g)(i)$ and $(\xi_n(f) \natural \xi_m(g))(i)$ equals to $n + m + 1 - f(i)$ or $m + 1 - g(i - n)$ depending whether $i \in \underline{n}$ or not. $\square$

An element $f \in \mathfrak{S}_n$ is called $\sharp$ -decomposable, provided $f = g \sharp h$ for some $g \in \mathfrak{S}_k, h \in \mathfrak{S}_m$. If such type decomposition is impossible, then f is called $\sharp$ -indecomposable. For each $f \in \mathfrak{S}_n$ we put

$$\delta(f) := \inf_i \{i \in \underline{n} \mid f(\underline{i}) \subset \underline{i}\}.$$

In particular $f(\{1, \dots, \delta(f)\}) \subset \{1, \dots, \delta f\}$. We let $f_\delta \in \mathfrak{S}_{\delta(f)}$ be the restriction of f on $\{1, \dots, \delta(f)\}$.

Lemma 3.5 i) If $f = g \sharp h$, then $\delta(f) = \delta(g)$ and $f_\delta = g_\delta$.

ii) If $f \in \mathfrak{S}_n$ is $\sharp$ -decomposable, then there is the unique $f_\beta \in \mathfrak{S}_{n-\delta(f)}$ such that

$$f = f_\delta \sharp f_\beta.$$

iii) An element $f \in \mathfrak{S}_n$ is $\sharp$ -indecomposable iff $f = f_\delta$.

Proof. i) It is sufficient to note that if $f = g \sharp h$ and $\mathcal{O}(g) = k$, then $f(\underline{k}) \subset f(\underline{k})$. To show ii) and iii) let us assume that $f = g \sharp h$. Then by i) we have

$$\delta(f) = \delta(g) \leq \mathcal{O}(g) < \mathcal{O}(f).$$

Thus $f \neq f_\delta$. Conversely, if $f \neq f_\delta$, then $f = f_\delta \sharp f_\beta$, where $f_\beta \in \mathfrak{S}_k$ is given by $f_\beta(i) = f(\delta(f) + i) - \delta(f)$. Here $k = \mathcal{O}(f) - \delta(f)$ and $1 \leq i \leq k$. The permutation f_β is well-defined since f maps the subset $\{\delta(f) + 1, \dots, \mathcal{O}(f)\}$ to itself. $\square$

We let $\mathfrak{U}^\sharp$ be the set of $\sharp$ -indecomposable elements of $\mathfrak{S}$. We have $\mathfrak{U}^\sharp = \coprod_n \mathfrak{U}_n^\sharp$, where $\mathfrak{U}_n^\sharp = \mathfrak{U}_n^\sharp \cap \mathfrak{S}_n$. For example we have

$$\mathfrak{U}_2^\sharp = \{(2, 1)\}, \quad \mathfrak{U}_3^\sharp = \{(2, 3, 1), (3, 1, 2), (3, 2, 1)\}$$

We let u_n be the cardinality of the set $\mathfrak{U}_n^\sharp$. Here are the first values of u_n :

$$1, 1, 3, 13, 71, 461, 3447, \dots$$

which can be deduced from Corollary 3.7. Jean-Louis Loday informed me that u_n is the number of permutations with no global descent [1].

Theorem 3.6 The semigroup $(\mathfrak{S}, \sharp)$ is free on $\mathfrak{U}^\sharp$.

Proof. First we show that the set $\mathfrak{U}^\sharp$ generates $\mathfrak{S}$. Take $f \in \mathfrak{S}$. We have to prove that f lies in the subsemigroup generated by $\mathfrak{U}^\sharp$. We may assume that $f \notin \mathfrak{U}^\sharp$. Thus f is $\sharp$ -decomposable and we can write $f = g \sharp h$. Since $\mathcal{O}(g), \mathcal{O}(h) < \mathcal{O}(f)$ we may assume by induction that g and h lie in the subsemigroup generated by $\mathfrak{U}^\sharp$. Thus the same is true for f as well. The fact that any f can be written uniquely in the form $f = g_1 \sharp \dots \sharp g_k$ with $g_i \in \mathfrak{U}^\sharp$ follows from Lemma 3.5. $\square$

Corollary 3.7 One has

$$\text{Card}(\mathfrak{U}_n^\sharp) = n! - \sum_{p+q=n} p!q! + \sum_{p+q+r=n} p!q!r! - \dots$$

Here $p, q, r, \dots$ runs over all strictly positive integers.

Proof. We have an isomorphism of graded sets $\mathfrak{S} \cong \coprod_{n \geq 1} \mathfrak{U}_n^\sharp$ and therefore

$$\sum_{n \geq 1} n! t^n = \sum_{n \geq 1} \left(\sum_{m \geq 1} \text{Card } \mathfrak{U}_m^\sharp t^m \right)^n$$

and the result follows. $\square$

By transportation of structures we see that $(\mathfrak{S}, \natural)$ is a free semigroup on the set $\mathfrak{U}^\natural$, which is by definition the set of all $\natural$ -indecomposable permutations. Here a permutation f is called $\natural$ -indecomposable if $\omega_n \circ f$ is $\sharp$ -indecomposable, in other words if $f(\{1, \dots, i\}) \not\subset \{n - i + 1, \dots, n\}$ for all $1 \leq i \leq n - 1$.

4 Free duplexes

4.1 Planar trees


By *tree* we mean in this paper a planar rooted tree. Such trees play an important rôle in the recent work of Jean-Louis Loday and Maria Ronco [6]. We let $\mathfrak{T}$ be the set of trees. It is a graded set

$$\mathfrak{T} = \coprod_{n \geq 1} \mathfrak{T}_n,$$

where $\mathfrak{T}_n$ is the set of trees with n leaves. The number of elements of $\mathfrak{T}_n$ are known as super Catalan numbers and they are denoted by C_n . Here are the first super Catalan numbers: $C_1 = 1 = C_2$, $C_3 = 3$, $C_4 = 11$, $C_5 = 45$. In general one has the following well-known relation (see for example Section 8.2 of [4]), which is also a consequence of Corollary 4.3.

$$f(\mathbf{T}) := \sum_{n \geq 1} C_n \mathbf{T}^n = \frac{1}{4} (\mathbf{T} + 1 - \sqrt{\mathbf{T}^2 - 6\mathbf{T} + 1}) \quad (1)$$

So $\mathfrak{T}_1$ has only one element $|$ and similarly $\mathfrak{T}_2$ has also only one element $\begin{array}{c} \diagup \quad \diagdown \\ | \end{array}$, while $\mathfrak{T}_3$ has three elements



Any vertex v of a tree $t \in \mathfrak{T}_n$ with $n \geq 2$, has a *level*, which is equal to the number of edges in the path connecting v to the root. Thus the root is the unique vertex of level 0. For example, if




then the tree u_1 has only one vertex (which is of course the root), while u_2 has two vertices the root and a vertex of level one.

Let us also recall that on trees there exists an important operation which is called *grafting*. The grafting defines a map

$$\mathbf{gr} : \mathfrak{T}_{n_1} \times \cdots \times \mathfrak{T}_{n_k} \rightarrow \mathfrak{T}_n, \quad n = n_1 + \cdots + n_k.$$

For example, if u_1 and u_2 are the same as above, then we have:



Let us also note, that $\mathbf{gr}(u_1, u_2)$ has 4 vertices, two of them of the level one and one of the level two. It is clear that any tree from $\mathfrak{T}_n$, $n > 1$ can be written uniquely as $\mathbf{gr}(t_1, \dots, t_k)$. Here k is the number of incoming edges at the root. We will say that t is constructed by the grafting of $(t_1, \dots, t_k)$.


We need also the following construction on trees. Let $t_1, \dots, t_k$ be trees and let I be a subset of $\underline{k}$. We consider $t = \mathbf{gr}(t_1, \dots, t_k)$. The tree obtained by contracting edges of t which connects the root of t with the roots of t_i , $i \in I$ is denoted by $\mathbf{gr}_I(t_1, t_2)$. If $I = \emptyset$, then $\mathbf{gr}_I = \mathbf{gr}$.

For example, if u_1 and u_2 are as above, then we have



4.2 The free duplex with one generator

We consider now two different copies of the set $\mathfrak{T}_n$ for $n \geq 2$, which are denoted by $\mathfrak{T}_n^\cdot$ and $\mathfrak{T}_n^*$, $n \geq 2$. We put $\mathfrak{T}^* := \bigcup_{n \geq 2} \mathfrak{T}_n^*$ and $\mathfrak{T}^\cdot := \bigcup_{n \geq 2} \mathfrak{T}_n^\cdot$. If $t \in \mathfrak{T}^\cdot$, then we assign $\cdot$ to any vertex of t of even level and $*$ to any vertex of t of odd level. Similarly, if $t \in \mathfrak{T}^*$, then we assign $*$ to any vertex of t of even level and $\cdot$ to any vertex of t of odd level. So, for example



We call such trees as decorated trees. To be more precise *a decorated tree* is an element of the set

$$\mathfrak{D} = \bigcup_{n=1}^{\infty} \mathfrak{D}(n),$$

where $\mathfrak{D}(1) = \mathfrak{T}_1$ and $\mathfrak{D}(n) = \mathfrak{T}_n \cup \mathfrak{T}_n^*$, $n \geq 2$. We are going to define a duplex structure on decorated trees in such a way, that the above tree can be expressed as

$$(e \cdot e \cdot e) * (e \cdot (e * e)),$$

where $e = |$. More formally, the operations

$$\cdot : \mathfrak{D} \times \mathfrak{D} \rightarrow \mathfrak{D}, \quad * : \mathfrak{D} \times \mathfrak{D} \rightarrow \mathfrak{D}$$

can be defined as follows. If $t_1, t_2 \in \mathfrak{T}_1$ then

$$t_1 * t_2 := \text{gr}(t_1, t_2) \in \mathfrak{T}_2^*, \quad t_1 \cdot t_2 := \text{gr}(t_1, t_2) \in \mathfrak{T}_2$$

if $t_1, t_2 \in \mathfrak{T}^*$ then

$$t_1 * t_2 := \text{gr}_{12}(t_1, t_2) \in \mathfrak{T}^*, \quad t_1 \cdot t_2 := \text{gr}(t_1, t_2) \in \mathfrak{T}$$

if $t_1, t_2 \in \mathfrak{T}$ then

$$t_1 * t_2 := \text{gr}(t_1, t_2) \in \mathfrak{T}^*, \quad t_1 \cdot t_2 := \text{gr}_{12}(t_1, t_2) \in \mathfrak{T}$$

if $t_1 \in \mathfrak{T}$ and $t_2 \in \mathfrak{T}^*$, then

$$t_1 * t_2 := \text{gr}_2(t_1, t_2) \in \mathfrak{T}^*, \quad t_1 \cdot t_2 := \text{gr}_1(t_1, t_2) \in \mathfrak{T}$$

if $t_1 \in \mathfrak{T}^*$ and $t_2 \in \mathfrak{T}$, then

$$t_1 * t_2 := \text{gr}_1(t_1, t_2) \in \mathfrak{T}^*, \quad t_1 \cdot t_2 := \text{gr}_2(t_1, t_2) \in \mathfrak{T}.$$

This concludes the construction of operations on $\mathfrak{D}$. Let us observe that in all cases

$$t_1 * t_2 = \text{gr}_I(t_1, t_2) \in \mathfrak{T}^*$$

and

$$t_1 \cdot t_2 = \text{gr}_J(t_1, t_2) \in \mathfrak{T},$$

where $I \subset \{1, 2\}$ consists with such $i \in \underline{2}$ that $t_i \in \mathfrak{T}^*$ and $J \subset \{1, 2\}$ consists with such $i \in \underline{2}$ that $t_i \in \mathfrak{T}$.

Lemma 4.1 *Both operations $*$ and $\cdot$ are associative.*

Proof. One observes that for any decorated trees t_1, t_2, t_3 in $\mathfrak{T}^*$ one has

$$(t_1 * t_2) * t_3 = \mathbf{gr}_I(t_1, t_2, t_3) = t_1 * (t_2 * t_3)$$

where $I \subset \{1, 2, 3\}$ consists with such i that $t_i \in \mathfrak{T}^*$, $i = 1, 2, 3$. Similarly for $\cdot$. $\square$

Lemma 4.2 *The semigroup $(\mathfrak{D}, *)$ is a free semigroup on the set*

$$S' := \mathfrak{T}_1 \cup \bigcup_{n \geq 1} \mathfrak{T}_n^* \subset \mathfrak{D}$$

and similarly, the semigroup $(\mathfrak{D}, \cdot)$ is a free semigroup on the set

$$S^* := \mathfrak{T}_1 \cup \bigcup_{n \geq 1} \mathfrak{T}_n \subset \mathfrak{D}$$

Proof. It suffices to note that if $t \in \mathfrak{D}$, but $t \notin S'$, then t can be written uniquely as $\mathbf{gr}(t_1, \dots, t_k)$. If we consider $t_1, \dots, t_k$ as elements from S' , then we have $t = t_1 * \dots * t_k$. Similarly for $(\mathfrak{D}, \cdot)$. $\square$

Comparing the corresponding formal power series we get the following well-known equation

Corollary 4.3 *For the super Catalan numbers we have*

$$\sum_{n \geq 1} (\mathbf{T} + \mathbf{T}^2 + 3\mathbf{T}^3 + 11\mathbf{T}^4 + \dots)^n = \mathbf{T} + 2(\mathbf{T}^2 + 3\mathbf{T}^3 + 11\mathbf{T}^4 + \dots).$$

Theorem 4.4 *The duplex $\mathfrak{D}$ is a free duplex with one generator $| \in \mathfrak{T}_1 \subset \mathfrak{D}$.*

Proof. For simplicity we let e be the tree $|$. Then e generates the duplex $\mathfrak{D}$. This can be proved by induction. Indeed, $\mathfrak{D}(2) = \mathfrak{T}_2 \cup \mathfrak{T}_2^*$ has two decorated trees, which are equal respectively to $e \cdot e$ and $e * e$. Assume we already proved that any decorated tree from $\mathfrak{D}(m)$ lies in the subduplex generated by a for any $m < n$ and let us prove that the same holds for $n = m$. Take $t \in \mathfrak{D}(n)$. Assume $t \in \mathfrak{T}_n^*$. Then there exists the unique trees $t_1, \dots, t_k$ such that $t = \mathbf{gr}(t_1, \dots, t_k)$. We consider $t_1, \dots, t_k$ as elements of the set $\mathfrak{T}_1 \cup \mathfrak{T}$. Then we have $t = t_1 * \dots * t_k$ and by the induction assumption t can be expressed in the terms of e . Similarly, if $t \in \mathfrak{T}$. Now we will show that $\mathfrak{D}$ is free on e . We take a duplex D and an element $a \in D$. We have to show that there exists a unique homomorphism $f : \mathfrak{D} \rightarrow D$ such that $f(e) = a$. We construct such f by induction. For the tree e we already have $f(e) = a$. Assume f is defined for all decorated trees from $\mathfrak{D}(m)$, $m < n$ and take a decorated tree $t \in \mathfrak{T}^n$. As above we can write $t = t_1 * \dots * t_k$

with unique $t_1, \dots, t_k$ and then we put $f(t) = f(t_1) * \dots * f(t_k)$. Similarly, if $t \in \mathfrak{T}$. Let us prove that f is a homomorphism. The equations

$$f(x * y) = f(x) * f(y), \quad f(x \cdot y) = f(x) \cdot f(y)$$

is clear if $x, y \in \mathfrak{T}_1$. Let us prove only the first one, because the proof of the second one is completely similar. If $x, y \in \mathfrak{T}$ then we have $x * y = \mathbf{gr}(x, y)$. Thus by definition $f(x * y) = f(x) * f(y)$. If $x \in \mathfrak{T}^*$ and $y \in \mathfrak{T}$. Then

$$x * y = \mathbf{gr}_1(x, y) = \mathbf{gr}(x_1, \dots, x_k, y) = x_1 * \dots * x_k * y$$

where $x = \mathbf{gr}(x_1, \dots, x_k)$ and $x_1, \dots, x_k$ are considered as elements of $\mathfrak{T}_1 \cup \mathfrak{T}$. It follows that $f(x * y) = f(x_1) * \dots * f(x_k) * f(y) = f(x) * f(y)$. Similarly if $x \in \mathfrak{T}$ and $y \in \mathfrak{T}^*$. If $x, y \in \mathfrak{T}^*$. Then

$$x * y = \mathbf{gr}_{12}(x, y) = \mathbf{gr}(x_1, \dots, x_k, y_1, \dots, y_l) = x_1 * \dots * x_k * y_1 * \dots * y_l$$

where $x = \mathbf{gr}(x_1, \dots, x_k)$, $y = \mathbf{gr}(y_1, \dots, y_l)$ and $x_1, \dots, x_k, y_1, \dots, y_l$ are considered as elements of $\mathfrak{T}_1 \cup \mathfrak{T}$. It follows that

$$f(x * y) = f(x_1) * \dots * f(x_k) * f(y_1) * \dots * f(y_l) = f(x) * f(y).$$

□

4.3 Free duplexes

For a set S we consider the set

$$\text{Dupl}(S) := \coprod_{n \geq 1} \text{Dupl}_n(S)$$

where $\text{Dupl}_n(S) = \mathfrak{D}(n) \times S^n$. We define maps

$$\cdot, * : \text{Dupl}_n(S) \times \text{Dupl}_m(S) \rightarrow \text{Dupl}_{n+m}(S)$$

by

$$(t_1, x_1, \dots, x_n) * (t_2, y_1, \dots, y_m) := (t_1 * t_2, x_1 \dots x_k, y_1 \dots y_m).$$

$$(t_1, x_1, \dots, x_n) \cdot (t_2, y_1, \dots, y_m) := (t_1 \cdot t_2, x_1 \dots x_k, y_1 \dots y_m).$$

We leave as an exercise to prove that $\text{Dupl}(S)$ is a free duplex on S . It is clear that for a finite set S one has

$$\text{Dupl}(S)(\mathbf{T}) = \mathbf{T} + 2 \sum_{n \geq 2}^{\infty} C_n (\text{Card}(S) \mathbf{T})^n. \quad (2)$$

5 Proof of Theorem 3.3

5.1 $\mathfrak{S}_2$ -indecomposable permutations

A permutation $f : \underline{n} \rightarrow \underline{n}$ is called $\mathfrak{S}_2$ -indecomposable if for any $i = 1, \dots, n-1$ one has

$$f(\underline{i}) \not\subset \underline{i}, \text{ and } f(\underline{i}) \not\subset \{n-i+1, \dots, n\}$$

It is clear that a permutation is $\mathfrak{S}_2$ -indecomposable iff it is simultaneously $\sharp$ - and $\natural$ -indecomposable and therefore for the set $\mathfrak{U}^{\mathfrak{S}_2}$ of $\mathfrak{S}_2$ -indecomposable permutations we have

$$\mathfrak{U}^{\mathfrak{S}_2} = \mathfrak{U}^{\sharp} \cap \mathfrak{U}^{\natural}. \quad (3)$$

It is clear that

$$\mathfrak{U}_1^{\mathfrak{S}_2} = \mathfrak{S}_1, \quad \mathfrak{U}_2^{\mathfrak{S}_2} = \emptyset = \mathfrak{U}_3^{\mathfrak{S}_2}, \quad \mathfrak{U}_4^{\mathfrak{S}_2} = \{(2, 4, 1, 3), (3, 1, 4, 2)\}$$

One checks that $\mathfrak{U}_5^{\mathfrak{S}_2}$ has 22 elements. In general for $d_n = \mathfrak{U}_n^{\mathfrak{S}_2}$ we have

$$d_n = n! - 2 \sum_{p+q=n} p!q! + 2 \sum_{p+q+r=n} p!q!r! - \dots = 2u_n - n! \quad (4)$$

Here $p, q, r, \dots$ runs over all strictly positive integers. This follows from the formula for $u_n = \text{Card} \mathfrak{U}_n^{\sharp}$ and from Lemma 5.1. Here are the first values of d_n :

$$1, 0, 0, 2, 22, 202, 1854, \dots$$

Lemma 5.1 *If a permutation is $\sharp$ -decomposable then it is $\natural$ -indecomposable. Similarly, if a permutation is $\natural$ -decomposable then it is $\sharp$ -indecomposable.*

Proof. If not, then there exists a permutation $f : \underline{n} \rightarrow \underline{n}$ and integers $1 \leq i \leq n-1$ and $1 \leq j \leq n-1$ with properties

$$f(\underline{i}) \subset \underline{i}, \text{ and } f(\underline{j}) \not\subset \{n-j+1, \dots, n\}.$$

Without loss of generality we may assume that $j \leq i$ (otherwise we take $\omega_n \circ f$). Thus for any $k = 1, \dots, j$ we have $n-j < f(k) \leq i$. Therefore the interval $[n-j, i]$ contains at least j integers. But this is impossible, because $i - n + j < j$. $\square$

5.2 Proof of Theorem 3.3

We will prove that $\mathfrak{S}$ is free on $\mathfrak{U}^{\mathfrak{S}_2}$ as a duplex. First we show that $\mathfrak{U}^{\mathfrak{S}_2}$ generates $\mathfrak{S}$. Indeed, if $f \notin \mathfrak{U}^{\mathfrak{S}_2}$ then either $f = g\sharp h$ or $f = g\natural h$ and it is impossible to have both cases. Without loss of generality we may assume that one has the

first possibility. Since $(\mathfrak{S}, \sharp)$ is a free semigroup, there exist uniquely defined $f_1, \dots, f_k \in \mathfrak{U}^\sharp$ such that $f = f_1 \sharp \dots \sharp f_k$. If all $f_i \in \mathfrak{U}^{\mathfrak{S}_2}$ we stop, otherwise for some i we have $f_i \notin \mathfrak{U}^\sharp$ and therefore one can write f_i in unique way as $f_i = g_1 \sharp \dots \sharp g_m$, with $g_j \in \mathfrak{U}^\sharp$. Since $\mathcal{O}(g_i) < \mathcal{O}(f)$ this process stops after a few steps. This shows that $\mathfrak{U}^{\mathfrak{S}_2}$ generates $\mathfrak{S}$ as a duplex. Since in each step there were unique choices we conclude that elements of $\mathfrak{U}^{\mathfrak{S}_2}$ are free generators. $\square$

Corollary 5.2 *Between the numbers d_n and super Catalan numbers C_n there is the following relation*

$$\sum_{n \geq 1} n! t^n = \sum_{n \geq 1} d_n t^n + 2 \sum_{m \geq 2} C_m \left(\sum_{n \geq 1} d_n t^n \right)^m$$

Furthermore one has

$$\psi^2 + \xi \psi - \psi + \xi = 0$$

where $\psi(\mathbf{T}) = \sum_{n \geq 1} n! \mathbf{T}^n$ and $\xi(\mathbf{T}) = \sum_{n \geq 1} d_n \mathbf{T}^n$.

Proof. The first part follows from Theorem 3.3 and Equation (2). To get the second part one rewrites the same relation as

$$\psi(\mathbf{T}) = 2f(\xi(\mathbf{T})) - \xi(\mathbf{T})$$

and then use the formula (1) for $f(\mathbf{T}) = \sum_{n \geq 1} C_n \mathbf{T}^n$. $\square$

6 Duplexes satisfying some additional identities

6.1 Duplex of binary trees

We let Duplexes_1 be the category of duplexes satisfying the identity

$$(a \cdot b) * c = a \cdot (b * c) \tag{5}$$

We prove that the free object in Duplexes_1 is given via binary trees. Let us recall that for us all trees are planar and rooted. A tree is called *binary* if any vertex is trivalent. We let $\mathfrak{Y}$ be the set of all binary trees:

$$\mathfrak{Y} := \coprod_{n \geq 1} \mathfrak{Y}_n,$$

where $\mathfrak{Y}_n$ is the set of planar binary trees with $n+1$ leaves. We have $\mathfrak{Y}_n \subset \mathfrak{T}_{n+1}$. The number of elements of $\mathfrak{Y}_n$ are known as Catalan numbers and they are

denoted by c_n . Here are the first Catalan numbers: $c_1 = 1$, $c_2 = 2$, $c_3 = 5$. In general one has

$$c_n = \frac{(2n)!}{n!(n+1)!}$$

So $\mathfrak{Y}_1$ has only one element $\begin{array}{c} \diagup \diagdown \\ | \end{array}$ while $\mathfrak{Y}_2$ has two elements



By our definition the tree $| \in \mathfrak{T}_1$ is not a binary tree, but it will play also an important rôle for binary trees as well. We let t be this particular tree. Let us note that any binary tree u can be written in the unique way as $\mathbf{gr}(u^l, u^r)$, where $u^l, u^r \in \mathfrak{Y} \cup \{t\}$. The map $\mathbf{gr} : \mathfrak{Y} \times \mathfrak{Y} \rightarrow \mathfrak{Y}$ is not anymore homogeneous, because we changed the grading, but it is of degree one: $\mathbf{gr} : \mathfrak{Y}_n \times \mathfrak{Y}_m \rightarrow \mathfrak{Y}_{n+m+1}$. We introduce two homogeneous operations

$$\cdot, * : \mathfrak{Y}_n \times \mathfrak{Y}_m \rightarrow \mathfrak{Y}_{n+m}$$

by

$$u \cdot v := \mathbf{gr}(u \cdot v^l, v^r)$$

$$u * v := \mathbf{gr}(u^l, u^r * v).$$

Here we use the induction and the convention $u \cdot t = t \cdot u = u * t = t * u = u$ for $t = |$.

These associative operations under the name 'over' and 'under' first appeared in [5]. One observes that (see [5]) $u \cdot v$ (resp. $u * v$) is the tree obtained by identifying the root of u (resp. v) with the left (resp. right) most leaf of v (resp. u).

Theorem 6.1 *The duplex $\mathfrak{Y}$ satisfies the identity (5). Moreover it is a free object in Duplexes_1 generated by $e = \mathbf{gr}(t, t) = \begin{array}{c} \diagup \diagdown \\ | \end{array}$.*

Proof. We have

$$(u \cdot v)^l = u \cdot v^l, \quad (u \cdot v)^r = v^r,$$

and

$$(u * v)^l = u^l, \quad (u * v)^r = u^r * v.$$

Therefore

$$(a \cdot b) * c = \mathbf{gr}(a \cdot b^l, b^r * c) = a \cdot (b * c).$$

Thus $\mathfrak{Y} \in \text{Duplexes}_1$. Now we prove that $\mathfrak{Y}$ is generated by e . First one observes that

$$u \cdot e = \mathbf{gr}(u, t), \quad e * u = \mathbf{gr}(t, u).$$

We claim that for any $a, b \in \mathfrak{Y} \cup \{t\}$ one has

$$(a \cdot e) * b = \mathbf{gr}(a, b)$$

Indeed, we can write $(a \cdot e) * b = \mathbf{gr}(a, t) * b$, which is the same as $\mathbf{gr}(a, t * b)$ thanks to the definition of the operation $*$. By our convention $t * b = b$ and the claim is proved. Take any element $u \in \mathfrak{Y}$. By the claim we have

$$u = (u^l \cdot e) * u^r = u^l \cdot (e * u^r) \quad (6)$$

Based on this equality one easily proves by the induction that e generates the duplex $\mathfrak{Y}$. Let D be any duplex satisfying the equality (5) and take any element $a \in D$. We have to show that there exists a unique homomorphism $f : \mathfrak{Y} \rightarrow D$ such that $f(e) = a$. Since e generates $\mathfrak{Y}$ the uniqueness is obvious. We construct recursively f by

$$f(e) = a, \quad \text{and} \quad f(u) = (f(u^l) \cdot a) * f(u^r)$$

and an obvious induction shows that f is indeed a homomorphism. $\square$

6.2 Dimonoids

We let **Dimonoids** be the full subcategory of **Duplexes**₁ satisfying two more identities:

$$(a * b) \cdot c = (a \cdot b) \cdot c \quad (7)$$

$$a * (b * c) = a * (b \cdot c) \quad (8)$$

We refer the reader to [2] for the extensive informations on dimonoids and *dialgebras* which are just linear analogs of dimonoids.

6.3 Duplex of vertices of cubes

We let **Duplexes**₂ be the full subcategory of **Duplexes**₁ satisfying the identity

$$(a * b) \cdot c = a * (b \cdot c) \quad (9)$$

Such type of algebras were first considered in [7] under the name “Doppelalgebren”. Free objects in **Duplexes**₁ are given via the following construction. For all $n \geq 2$ we let $\mathfrak{Q}_n$ to be the set of vertices of $n - 1$ dimensional cube. So $\mathfrak{Q}_n = \{-1, 1\}^{n-1}$ and elements of $\mathfrak{Q}_n$ are sequences $\mathbf{a} = (a_1 \cdots, a_{n-1})$, where $a_i = -1$ or $a_i = 1$. Moreover we let $\mathfrak{Q}_1$ to be the singleton $\{e\}$ and

$$\mathfrak{Q} := \coprod_{n \geq 1} \mathfrak{Q}_n.$$

Following to [5] we define two homogeneous operations on $\mathfrak{Q}$ by

$$\begin{aligned}
e \cdot e &:= -1 \in \mathfrak{Q}_2, \\
e * e &= 1 \in \mathfrak{Q}_2, \\
e \cdot \mathbf{a} &:= (-1, a_1, \dots, a_{n-1}), \\
\mathbf{a} \cdot e &:= (a_1, \dots, a_{n-1}, -1), \\
e * \mathbf{a} &:= (1, a_1, \dots, a_{n-1}), \\
\mathbf{a} * e &:= (a_1, \dots, a_{n-1}, 1), \\
\mathbf{a} \cdot \mathbf{b} &:= (a_1, \dots, a_{n-1}, -1, b_1, \dots, b_{m-1}), \\
\mathbf{a} * \mathbf{b} &:= (a_1, \dots, a_{n-1}, 1, b_1, \dots, b_{m-1}).
\end{aligned}$$

One checks that $\mathfrak{Q}$ is a duplex, which satisfy both identities (5) and (9).

Theorem 6.2 [7] *The duplex $\mathfrak{Q}$ is the free object in $\mathbf{Duplexes}_2$ generated by e .*

Proof. We first show that e generates $\mathfrak{Q}$. For $\mathbf{a} = (a_1, \dots, a_n) \in \mathfrak{Q}_n$, we put $\circ_i = \cdot$ if $a_i = -1$ and $\circ_i = *$ if $a_i = 1$. Then $\mathbf{a} = e \circ_1 e \circ_2 \dots \circ_{n-1} e$. The fact that this expression does not depends on parenthesis follows from the associativity of $\cdot, *$ and from the identities (5),(9). Let D be an object of $\mathbf{Duplexes}_2$ and let $x \in D$. One can use the expression of a in terms of e and $\circ_i$ to define the map $f : \mathfrak{Q} \rightarrow D$ by $f(\mathbf{a}) = x \circ_1 x \circ_2 \dots \circ_{n-1} x$. Then by induction one shows that f is in fact a homomorphism with $f(e) = x$. $\square$

6.4 Remarks

Our results can be used to describe maps between combinatorial objects. It is not difficult to check that the map $\psi : \mathfrak{S} \rightarrow \mathfrak{Y}$ constructed in [5] is a homomorphism of duplexes. On the other hand, since $\mathfrak{D}$ is a free duplex with one generator there is a unique homogeneous map $\alpha : \mathfrak{D} \rightarrow \mathfrak{S}$ which is also a homomorphism of duplexes. To specify this map it suffice to know the image of the generator: $\alpha(|) = Id_1$. Similarly, if one considers $\mathfrak{Y}$ as an object of $\mathbf{Duplexes}$, then there exist a unique surjective homomorphism $\varrho : \mathfrak{D} \rightarrow \mathfrak{Y}$ which takes the generator $|$ of $\mathfrak{D}$ to $e = \mathbf{gr}(t, t) \in \mathfrak{Y}_1$. Thus we have $\varrho = \psi \circ \alpha$. Since $\mathbf{Duplexes}_2 \subset \mathbf{Duplexes}_1$, the free object on a set X in $\mathbf{Duplexes}_2$ is a quotient of the free object on a set X in $\mathbf{Duplexes}_1$. In particular for $X = \{e, \}$ we obtain the canonical quotient homomorphism $\phi : \mathfrak{Y} \rightarrow \mathfrak{Q}$. If one forgets the corresponding algebraic structures this map from binary trees to vertices of cubes coincides with one considered in [5]. The composite map $\phi \circ \varrho : \mathfrak{D} \rightarrow \mathfrak{Q}$ from the decorated trees to the vertices has the following description. Take a decorated tree $u \in \mathfrak{D}_n$ and label the leaves of u by the numbers $1, \dots, n$. Then $\phi \circ \varrho(u) = (a_1, \dots, a_{n-1})$, where a_i is $+1$ if

the sign at the end of the edge coming from the $(i + 1)$ -th leaf is $*$ and is -1 if the corresponding sign is $.$

Acknowledgments

This paper is influenced by the work of Jean-Louis Loday, who started to investigate algebras with two associative operations (see [2, 3]) and initiated me to the beauty of the combinatorics of trees during several informal discussions. This work was written during my visit at Universität Bielefeld. I would like to thank Friedhelm Waldhausen for the invitation to Bielefeld. Thanks to my wife Tamar Kandaorishvili for various helpful discussions on the subject and encouraging me to write this paper. The author was partially supported by the grant INTAS-99-0081 and TMR Network ERB FMRX CT-97-0107.

References

- [1] M. AGUIAR and F. SOTTILE. *Structures of the Malvenuto-Reutenauer Hopf algebra of permutations*. arXiv:math.CO/0203282.
- [2] J.-L. LODAY. *Algèbres ayant deux opérations associatives (digèbres)*. C. R. Acad. Sci. Paris Sér. I Math. 321 (1995), no. 2, 141–146.
- [3] J.-L. LODAY. *Dialgebras*. in "Dialgebras and related operads", Lecture Notes in Math., 1763, Springer, Berlin, (2001), 7–66.
- [4] J.-L. LODAY. *Arithmetree*. J. of Algebra. 258(2002), 275-309.
- [5] J.-L. LODAY and M. O. RONCO. *Order structure on the algebra of permutations and of planar binary trees*. J. Algebraic Combin. 15 (2002), no. 3, 253–270
- [6] J.-L. LODAY and M. O. RONCO. *Tri-algebras and families of polytopes*. Preprint, math.AT/0205043
- [7] B. RICHTER. *Dialgebren, Doppelalgebren und ihre Homologie*. Diplomarbeit. Universität Bonn. 1997. [www.math.uni-bonn.de/people/richter/]